Research on Standardization and Risk Control of Clinical Operations in Urology from the Perspective of Medical Technology Innovation

Lei Xing[1], Xin Zhao[2]

[1,2] *The Second People's Hospital of Kunming, Kunming 650000, China*
[1] *Email: 872762704@qq.com*
[2] *Corresponding author. Email: 875537592@qq.com*

ABSTRACT
This article focuses on the important issues of clinical operation of urology under the wave of medical technology innovation. With the continuous innovation of medical technology, the clinical operation of urology is facing new challenges of standardization and risk control. In this paper, a large number of related materials are sorted out by literature research, and the internal relationship between medical technology innovation and clinical operation norms and risk control of urology is deeply analyzed by logical analysis. It is found that the innovation of medical technology not only brings opportunities to the clinical operation of urology, but also causes problems such as lagging operation norms and various types of risks. In view of these situations, this paper proposes to build a standardization strategy from two aspects: norm formulation and promotion, covering measures such as multidisciplinary collaboration in norm formulation, dynamic updating of norm content, and diversified training and information platform promotion. In this paper, the risk control strategy is implemented from three stages: preoperative, intraoperative and postoperative, including building a comprehensive evaluation system, real-time monitoring and emergency treatment of advanced equipment, and improving follow-up and data processing. These strategies are helpful to improve the medical quality of urology and ensure the safety of patients.

Keywords: *Medical technology innovation, Urology, Standardization of clinical operation, Risk control, Evidence-based medicine.*

1. INTRODUCTION

The rapid development of medical technology has brought unprecedented changes to modern medicine, and the field of urology has also benefited greatly [1]. In recent years, an endless stream of innovative medical technology, from advanced diagnostic equipment to complex surgical operation technology, has greatly changed the diagnosis and treatment mode of urological diseases [2]. These technological innovations not only provide doctors with more accurate and efficient diagnosis and treatment methods, but also bring more hope for patients to be cured [3]. While enjoying the bonus of technological innovation, the clinical operation of urology is facing new challenges. The standardization of clinical operation is directly related to medical quality and patient safety. Under the background of rapid upgrading of medical technology, traditional clinical operation norms need to keep pace with the times [4]. If the new technology cannot be incorporated into the norms in time, it may lead to confusion and disorder in clinical operation and increase medical risks.

The clinical operation of urology is highly professional and complex, involving the fine structure and important functions of human urinary system. A small operational error may bring serious consequences to patients [5]. Therefore, ensuring the standardization of clinical operation is always the core requirement of urology medical practice [6]. Risk control is an important link to ensure medical safety, which runs through the whole

process of clinical operation, from preoperative evaluation to postoperative rehabilitation.

It is of great practical significance to study the standardization of clinical operation and risk control in urology. This is helpful to improve the overall medical level of urology and promote the healthy development of the discipline [7]. By clarifying the operating norms under the background of new technology, doctors can use various medical technologies more scientifically and reasonably, and improve the treatment effect [8]. This can also effectively protect the rights and interests of patients and reduce medical risks. Standardized operation and effective risk control can reduce the occurrence of complications, improve the quality of patients' rehabilitation and enhance their trust in medical services. The purpose of this study is to deeply analyze the related problems of standardization of clinical operation and risk control in urology from the unique perspective of medical technology innovation. By systematically sorting out the specific manifestations of medical technology innovation in urology, this paper analyzes its influence on clinical operation norms, identifies potential risk types and explores the causes. Based on this, the targeted standardized strategies and risk control measures are put forward to provide support for improving the medical quality of urology.

2. INFLUENCE OF MEDICAL TECHNOLOGY INNOVATION ON STANDARDIZATION OF CLINICAL OPERATION IN UROLOGY DEPARTMENT

The innovation of medical technology not only brings great progress to urology, but also affects the standardization of clinical operation in many aspects, which brings both opportunities and challenges. From the perspective of opportunity, technological innovation provides a strong support for the standardization of clinical operation. New diagnostic techniques, such as high-resolution imaging equipment, can show the details of urinary system lesions more clearly, so that doctors have a more accurate basis in the diagnosis process, which provides a solid foundation for formulating standardized diagnostic procedures [9]. In terms of treatment, the development of minimally invasive technology and robot-assisted surgery, because of its advantages of less trauma and quick recovery, has promoted the clinical operation to a more refined and standardized direction.

The innovation of medical technology has also brought many challenges to the standardization of clinical operation. At the initial stage of new technology application, there is often a lag in operating norms. Because the period from research and development to clinical application of new technology is short, the relevant operating specifications may not be perfect synchronously, and doctors lack clear guidance when using new technology, which is prone to irregular operation [10]. The exploration and application of some cutting-edge urological techniques, such as gene editing technology in the treatment of urinary system diseases, involves multidisciplinary knowledge, and the formulation of its operational norms requires the cooperation of interdisciplinary experts. The standardization and promotion of these complex technologies are also faced with the limitations of knowledge updating speed and training resources of medical staff, which makes it difficult to implement the standardization in clinical practice quickly and effectively.

3. RISK TYPES AND CAUSES OF CLINICAL OPERATION IN UROLOGY DEPARTMENT

3.1 Risk Types

Surgical operation risk is common in clinical operation of urology. In kidney surgery, there may be massive bleeding during operation. This is mainly due to the rich blood supply of the kidney. Once the large blood vessels are damaged during the operation, it will cause serious bleeding, which will affect the operation process and the safety of patients. Prostate surgery may lead to urethral sphincter injury, and then lead to urinary incontinence, which has a great negative impact on the quality of life of patients after operation.

The risk of postoperative complications is also an important aspect of urology clinic. Take urinary system infection as an example. After operation, patients' body resistance decreases, and indwelling catheter and other factors can easily lead to bacterial growth and infection. Urinary fistula is also one of the common postoperative complications, which is mostly caused by poor healing of the surgical site and leakage of urine to surrounding tissues, which not only increases the pain of patients, but also may lead to more serious infection.

The risk of diagnosis error cannot be ignored. On the one hand, misdiagnosis may occur, such as

misdiagnosis of urinary calculi as other abdominal diseases, delaying the treatment opportunity. On the other hand, missed diagnosis also occurs from time to time. When the early symptoms of some concealed urinary system tumors are not obvious, they may be missed because of incomplete examination and miss the best treatment period.

3.2 Cause Analysis

With the rapid development of medical technology, some new treatment methods and equipment are constantly applied in urology clinic. However, in the initial stage of application of new technology, doctors are not fully aware of its operation skills and potential risks. The application of new technology may change the natural course and clinical manifestations of the disease, making the traditional diagnostic criteria and methods no longer fully applicable and increasing the risk of diagnostic errors. “Figure 1” shows the risk types and causes of clinical operation in urology:




Figure 1 Types and causes of clinical operation risks in urology.

Some doctors may violate the established operating procedures during the operation, such as not strictly following the principle of aseptic operation, which increases the risk of postoperative infection. Inexperienced doctors may also lead to operational errors. Because young doctors have relatively little practical experience, they may not be able to make accurate judgments and respond effectively when dealing with complex diseases or emergencies.

4. STANDARDIZATION STRATEGY OF CLINICAL OPERATION OF UROLOGY BASED ON MEDICAL TECHNOLOGY INNOVATION

4.1 Specification Formulation Strategy

Medical technology innovation often involves many disciplines, so it is the key to establish a multi-disciplinary collaborative norm-setting team. The team should include urologists, medical engineers, biologists and evidence-based medicine experts. With rich clinical experience, urologists can put forward demands from the practical point of view. Medical engineers can provide technical suggestions for operating specifications according to technical principles. Biologists can start from the basic theory of life science and assist in optimizing norms. Evidence-based medicine experts are responsible for integrating all kinds of research evidence to ensure that the norms are scientific and reliable. Through the in-depth cooperation of multi-disciplines, the operation specification not only fits the clinical practice, but also conforms to the technical principle and scientific basis. Medical technology innovation is changing with each passing day, so the clinical operation norms of urology must keep up with the pace of technical development, establish a dynamic updating mechanism, and regularly review and revise the norms.

4.2 Standardization Promotion Strategy

Developing diversified training activities is an important means to popularize clinical practice norms. Hospitals can organize offline centralized training courses and invite authoritative experts in the industry to give theoretical explanations and operational demonstrations, so that medical staff can intuitively understand the specific contents and operational points of the norms. Use online learning platform to provide rich teaching resources, such as operation videos and case studies, to facilitate medical staff to learn anytime and anywhere. “Figure 2” shows the standardization strategy of urology clinical operation based on medical technology innovation:




Figure 2 Standardization strategies for clinical operations in urology.

With the help of information platform, it can realize the convenient query and real-time update of clinical operation specifications. In the information platform, it is necessary to establish a special database of clinical operation norms of urology, and systematically sort out and classify all kinds of norms. Medical staff can quickly query the required operating specifications through the hospital internal network or mobile terminal. When the specification content is updated, the system can push the notification in time to ensure that the medical staff use the latest version of the specification. The information platform can also be equipped with interactive communication function, and medical staff can discuss and consult on the platform at any time when they encounter problems in practice, so as to promote the effective promotion and application of norms.

5. RISK CONTROL STRATEGY OF UROLOGY CLINICAL OPERATION BASED ON MEDICAL TECHNOLOGY INNOVATION

Under the background of medical technology innovation, the clinical operation of urology is facing new risk challenges. In order to ensure the safety of patients and improve the quality of medical care, it is very important to implement effective risk control strategies. “Figure 3” presents the specific contents of the risk control strategy at each stage of urology clinical operation based on medical technology innovation:




Figure 3 Risk control strategies for clinical operations in urology.

5.1 Preoperative Risk Assessment

Comprehensive and accurate preoperative risk assessment is the first step of risk control. Doctors should comprehensively consider the patient’s individual condition, disease characteristics and the characteristics of innovative medical technology to be adopted. For example, for patients who plan to undergo a new robot-assisted urological tumor resection, it is necessary not only to evaluate their basic physical conditions such as cardiopulmonary function and coagulation mechanism, but also to consider the influence of this new technology on the location, size and surrounding tissue relationship of the patient’s specific tumor. Therefore, doctors should build a complete preoperative risk assessment system, including detailed patient history collection, comprehensive physical examination and special evaluation indicators for innovative technologies.

5.2 Intraoperative Risk Monitoring

Intraoperative risk monitoring is the key to ensure the smooth operation. With the help of advanced monitoring equipment, doctors can track the patient's vital signs, important organ functions and key indicators of the surgical site in real time. For example, when the new laser lithotripsy technology is used, the temperature change of local tissues in the process of lithotripsy is monitored in real time through special optical fiber sensors to prevent thermal damage. The surgical team should have a high degree of emergency handling ability and make detailed plans for possible sudden risks.

5.3 Postoperative Risk Tracking

Postoperative risk tracking cannot be ignored. Medical staff should establish a perfect postoperative follow-up system and pay close attention to the rehabilitation process of patients. Medical staff regularly checks up patients, ask about the changes of symptoms, and find and deal with potential risks after operation in time. For patients treated with innovative technologies, special attention should be paid to the special complications that new technologies may cause. Systematic collection and in-depth analysis of postoperative risk data, summing up experience and lessons, and providing valuable reference for risk control of similar operations in the future.

6. CONCLUSION

This study focuses on the standardization of clinical operation and risk control of urology from the perspective of medical technology innovation. The innovation of medical technology has profoundly changed the pattern of clinical operation in urology, which not only brings opportunities such as accurate diagnosis and efficient treatment, but also causes many challenges in operational norms and risk prevention and control. In terms of standardization, the formulation of norms should rely on multidisciplinary cooperation and integrate the professional knowledge of all parties to ensure that the norms are in line with clinical practice and have scientific basis, and should be updated in time according to the development of technology. Standardization promotion should combine diversified training with information platform to effectively improve medical staff's cognition and implementation ability of standardization. In terms of risk control, a comprehensive risk assessment system before operation can comprehensively consider the individual patients and new technical factors, and provide a reliable basis for the surgical plan. Real-time monitoring of advanced equipment during operation and perfect emergency handling mechanism can effectively deal with all kinds of sudden risks. The perfect follow-up system and systematic data processing after operation are helpful to find and deal with potential risks in time and continuously optimize risk control strategies.

Only by fully implementing the strategy of clinical operation standardization and risk control can urology make steady progress on the road of medical technology innovation, effectively improve medical quality, protect patients' rights and interests, and lay a solid foundation for the development of urology.

ACKNOWLEDGEMENTS

Kunming Health Science and Technology Talent Development Project – Training Plan for Leaders in Medical Science and Technology, Project Number: 2023-SW(Leader)-26.

REFERENCES

[1] Dong Jinkai, Guo Wei, Hu Jinchao, et al. Risk Management in Various Stages of Clinical Trials in Urology: Summary of Single-Center Experience[J]. Journal of Modern Urology, 2020, 25(09): 765-767+783. DOI: 10.3969/j.issn.1009-8291.2020.09.001.

[2] Huang Helou, Gao Jianping, Sun Qing, et al. Research on the Value of a Dynamic Feedback and Control System in the Operation Management of Urological Surgical Equipment[J]. China Medical Equipment, 2022, 19(08): 149-153. DOI: 10.3969/J.ISSN.1672-8270.2022.08.032.

[3] Bai Xiaojing, Xie Na, Ma Yafang, et al. Evaluation and Effect Analysis of Urological Equipment Operation Quality Based on the Pressure-State-Response Model[J]. China Medical Equipment, 2025, 22(05): 93-98.

[4] Zhang Jingyun, Fu Chengwei, Tian Weilan, et al. Effect of Nursing Management Intervention Model Based on Infection Control on Preventing Adverse Events in Urology[J]. Chinese Journal of Disinfection, 2021, 38(12): 929-931. DOI: 10.11726/j.issn.1001-7658.2021.12.015.

[5] Liang Qianyan, Huang Liufang, Wu Rong, et al. Observation on the Effect of 90° Lateral Oblique Position in Retroperitoneal Laparoscopic Urological Surgery[J]. Modern Clinical Nursing, 2024, 23(07): 23-28. DOI: 10.3969/j.issn.1671-8283.2024.07.004.

[6] Meng Xianli, Cao Jie, Cheng Xin, et al. A Qualitative Study on the Psychological Feelings of Urology Nursing Interns Performing Care Operations for Patients in Private Areas[J]. Chinese Nursing Management, 2022, 22(04): 533-536.

[7] Yan Fengqi, Jiao Yong, Cui Dong, et al. Application Effect of 3D Printing Technology in Clinical Teaching of Percutaneous Nephrolithotomy in Standardized Residency Training[J]. China Medical Herald, 2022, 19(22): 71-74. DOI: 10.20047/j.issn1673-7210.2022.22.016.

[8] Dong Dexin, Wang Wenda. Application of Peyton's Four-Step Teaching Method in Clinical Internship Teaching in Urology[J]. Basic & Clinical Medicine, 2024, 44(07): 1058-1060. DOI: 10.16352/j.issn.1001-6325.2024.07.1058.

[9] Zhang Ying, Zhang Yanli, Mi Xue, et al. Exploration of the Application of a Performance Appraisal Program in Urology Nursing Management[J]. Journal of Nurses Training, 2020, 35(20): 1883-1886. DOI: 10.16821/j.cnki.hsjx.2020.20.008.

[10] Zhang Yan, Xu Lifen, Luo Jian, et al. Design and Implementation of Clinical Teaching in Humanistic Care for Urology Nursing Interns[J]. Journal of Nursing Science, 2021, 36(17): 75-77. DOI: 10.3870/j.issn.1001-4152.2021.17.075.